

**Borshchevska A.V.,**  
*Candidate of Medical Sciences,  
Senior Lecturer psychiatry and psychotherapy department,  
Faculty of postgraduate education,  
Lviv National Medical University named after Danylo Halytskyi*

## **BULLING AT THE SCHOOLS AS A FACTOR OF MENTAL ILL HEALTH**

Children and young people are the most sensitive group as to changes of any kind, which as a calque reflects modifications taking place in the modern society. Two main important systems, which take place in the child development are family and pedagogic surroundings. Unhealthy atmosphere of some schools gives possibility to thriving of the violence development, to be more precise-persecution aggression, the term for which is “bullying”. The latter influences negatively the possibility to study successfully, their psychological and psychical state.

The leading countries that education is a social lift that allows to improve the life chances of every individual, regardless of social status and place of residence. “School – is the main area, which is the practice of growing. The results showed that it is the ratio of the head of the educational institution to the classroom is a leading factor in determining social and psychological climate of an educational institution, regardless of the socioeconomic level of the school. All students 8-10 years are able to use tools bully, although about 40% of children use it often. At the same time some of the children – about 13% – are actively using baiting. Most prevalent (about 40%) with

verbal abuse – the most safe and those that impunity for aggressors, in second place (about 20% each) are physical abuse and mental oppression, at the Place of the latter (about 15%) – the prohibition and neglect (stigma). In the transition from elementary school to high bully mostly weak. But of some children – a stable personal trait, with age only change its form of manifestation. It is clear that this feature is “born” in the family and carries the imprints of the specifics of the family system. As for family risk factors bully then they can be considered: the absence of his father, depressed mother, violence in the family. Children get used to behave aggressively in relation to the weak, and observing the daily life of the family. Often buleramy are children whose parents teach dominant treatment, suppression of others, defending their interests. Children who suffer from parental cruelty and psychopathy, abandoned, deprived of maternal warmth and tenderness, as often became ahressors. Medicine, including child psychiatry, may become the moderator, who will help establish a common language for all adults and organize meaningful opportunity to be heard and to get timely help a child that is in stressful situation harassment.